arXiv:1605.05610v2 [cs.NA] 20 May 2016

A Short Proof for Gap Independence of Simultaneous Iteration

Edo Liberty
Yahoo Research

This note provides a very short proof of a spectral gap independent property of the simultaneous iterations algorithm for finding the top singular space of a matrix [1, 2, 3, 4]. The proof is terse but completely self contained and should be accessible to the linear algebra savvy reader.

Lemma. *Let $A \in \mathbb{R}^{n\times m}$ be an arbitrary matrix and let $G \in \mathbb{R}^{m\times k}$ be a matrix of i.i.d. random Gaussian entries. Let $t = c \cdot \log(n/\varepsilon)/\varepsilon$ and $Z = \text{span}((AA^T)^t AG)$ then with high probability depending only on a universal constant c*

$$||A - ZZ^T A|| \leq (1+\varepsilon)\sigma_{k+1}$$

Proof. $||A - ZZ^T A|| = \max_{x:\|x\|=1} \|x^T A\|$ such that $\|x^T Z\| = 0$. Using the SVD of A we change variables $A = USV^T$, $x = Uy$ and $G' = V^T G$. Note that G' is also a matrix of i.id. Gaussian entries because V is orthogonal. We get $||A - ZZ^T A|| = \max_{y:\|y\|=1} \|y^T S\|$ s.t. $y^T S^{2t+1} G' = 0$. We now break y, S, and G' to two blocks each such that

$$y = \left(\frac{y_1}{y_2}\right), \; S = \left(\begin{array}{c|c} S_1 & 0 \\ \hline 0 & S_2 \end{array}\right), \; G' = \left(\frac{G'_1}{G'_2}\right)$$

and $y_1 \in \mathbb{R}^k$, $y_2 \in \mathbb{R}^{n-k}$, $S_1 \in \mathbb{R}^{k\times k}$, $S_2 \in \mathbb{R}^{(n-k)\times(n-k)}$, $G'_1 \in \mathbb{R}^{k\times k}$, and $G'_2 \in \mathbb{R}^{(n-k)\times k}$.

$$\begin{aligned} 0 &= \|y^T S^{2t+1} G'\| = \|y_1^T S_1^{2t+1} G'_1 + y_2^T S_2^{2t+1} G'_2\| \\ &\geq \|y_1^T S_1^{2t+1} G'_1\| - \|y_2^T S_2^{2t+1} G'_2\| \\ &\geq \|y_1^T S_1^{2t+1}\| / \|G'^{-1}_1\| - \|y_2^T\| \cdot \|S_2^{2t+1}\| \cdot \|G'_2\| \\ &\geq |y_1(i)| \sigma_i^{2t+1} / \|G'^{-1}_1\| - \sigma_{k+1}^{2t+1} \cdot \|G'_2\| \; . \end{aligned}$$

This gives that $|y_1(i)| \leq (\sigma_{k+1}/\sigma_i)^{2t+1} \|G'_2\| \|G'^{-1}_1\|$. Equipped with this inequality we bound the expression $\|y^T S\|$. Let $1 \leq k' \leq k$ be such that $\sigma_{k'} \geq (1+\varepsilon)\sigma_{k+1}$ and $\sigma_{k'+1} < (1+\varepsilon)\sigma_{k+1}$. If no such k' exists the claim is trivial.

$$\begin{aligned} ||A - ZZ^T A||^2 &= \|y^T S\|^2 = \sum_{i=1}^{k'} y_i^2 \sigma_i^2 + \sum_{i=k'+1}^{n} y_i^2 \sigma_i^2 & (1) \\ &\leq (\|G'_2\|^2 \|G'^{-1}_1\|^2 \sum_{i=1}^{k'} (\sigma_{k+1}/\sigma_i)^{4t} \sigma_{k+1}^2) + (1+\varepsilon)\sigma_{k+1}^2 & (2) \\ &\leq \left[\|G'_2\|^2 \|G'^{-1}_1\|^2 k (1/(1+\varepsilon))^{4t} + (1+\varepsilon)\right] \sigma_{k+1}^2 \leq (1+2\varepsilon)\sigma_{k+1}^2 & (3) \end{aligned}$$

The last step is correct as long as $\|G'_2\|^2 \|G'^{-1}_1\|^2 k (1/(1+\varepsilon))^{4t} \leq \varepsilon \sigma_{k+1}^2$ which holds for $t \geq \log(\|G'_2\|^2 \|G'^{-1}_1\|^2 k/\varepsilon)/4\log(1+\varepsilon) = O(\log(n/\varepsilon)/\varepsilon)$. The last inequality uses the fact that G'_1 and G'_2 are random gaussian due to rotational invariance of the Gaussian distribution. This means that $\|G'_2\|^2 \|G'^{-1}_1\|^2 = O(\text{poly}(n))$ with high probability [5]. Finally, $||A - ZZ^T A|| \leq \sqrt{1+2\varepsilon} \cdot \sigma_{k+1} \leq (1+\varepsilon)\sigma_{k+1}$. ∎

References

[1] Vladimir Rokhlin, Arthur Szlam, and Mark Tygert. A randomized algorithm for principal component analysis. *SIAM J. Matrix Analysis Applications*, 31(3):1100–1124, 2009.

[2] N. Halko, P. G. Martinsson, and J. A. Tropp. Finding structure with randomness: Probabilistic algorithms for constructing approximate matrix decompositions. *SIAM Rev.*, 53(2):217–288, May 2011.

[3] Cameron Musco and Christopher Musco. Randomized block krylov methods for stronger and faster approximate singular value decomposition. In Corinna Cortes, Neil D. Lawrence, Daniel D. Lee, Masashi Sugiyama, and Roman Garnett, editors, *Advances in Neural Information Processing Systems 28: Annual Conference on Neural Information Processing Systems 2015, December 7-12, 2015, Montreal, Quebec, Canada*, pages 1396–1404, 2015.

[4] Rafi Witten and Emmanuel Candès. Randomized algorithms for low-rank matrix factorizations: Sharp performance bounds. *Algorithmica*, 72(1):264–281, May 2015.

[5] Mark Rudelson. Invertibility of random matrices: norm of the inverse. *Annals of Mathematics*, 168 Issue 2:575–600, 2008.